

PROOF COVER SHEET

Author(s): Vito Lubes

Article title: Mixed-ligand complex formation equilibria of vanadium(III) with picolinic and dipicolinic acids with some dicarboxylic acids (oxalic, malonic and phthalic acids) studied in 3.0 M KCl at 25 °C

Article no: TCSB 1023087

Enclosures: 1) Query sheet
2) Article proofs

Dear Author,

1. Please check these proofs carefully. It is the responsibility of the corresponding author to check these and approve or amend them. A second proof is not normally provided. Taylor & Francis cannot be held responsible for uncorrected errors, even if introduced during the production process. Once your corrections have been added to the article, it will be considered ready for publication.

Please limit changes at this stage to the correction of errors. You should not make trivial changes, improve prose style, add new material, or delete existing material at this stage. You may be charged if your corrections are excessive (we would not expect corrections to exceed 30 changes).

For detailed guidance on how to check your proofs, please paste this address into a new browser window:
<http://journalauthors.tandf.co.uk/production/checkingproofs.asp>

Your PDF proof file has been enabled so that you can comment on the proof directly using Adobe Acrobat. If you wish to do this, please save the file to your hard disk first. For further information on marking corrections using Acrobat, please paste this address into a new browser window: <http://journalauthors.tandf.co.uk/production/acrobat.asp>

2. Please review the table of contributors below and confirm that the first and last names are structured correctly and that the authors are listed in the correct order of contribution. This check is to ensure that your name will appear correctly online and when the article is indexed.

Sequence	Prefix	Given name(s)	Surname	Suffix
1		Ana	Esteves	
2		Jessaid	Escobar	
3		José Daniel	Martínez	
4		Mary Lorena	Araujo	
5		Felipe	Brito	
6		Lino	Hernández	
7		Edgar	Del Carpio	
8		Vito	Lubes	

Queries are marked in the margins of the proofs, and you can also click the hyperlinks below. Content changes made during copy-editing are shown as tracked changes. Inserted text is in **red font** and revisions have a **red** indicator **^**. Changes can also be viewed using the list comments function. To correct the proofs, you should insert or delete text following the instructions below, but **do not add comments to the existing tracked changes**.

AUTHOR QUERIES

General points:

1. **Permissions:** You have warranted that you have secured the necessary written permission from the appropriate copyright owner for the reproduction of any text, illustration, or other material in your article. Please see <http://journalauthors.tandf.co.uk/permissions/usingThirdPartyMaterial.asp>.
2. **Third-party content:** If there is third-party content in your article, please check that the rightsholder details for re-use are shown correctly.
3. **Affiliation:** The corresponding author is responsible for ensuring that address and email details are correct for all the co-authors. Affiliations given in the article should be the affiliation at the time the research was conducted. Please see <http://journalauthors.tandf.co.uk/preparation/writing.asp>.
4. **Funding:** Was your research for this article funded by a funding agency? If so, please insert ‘This work was supported by <insert the name of the funding agency in full>’, followed by the grant number in square brackets ‘[grant number xxxx]’.
5. **Supplemental data and underlying research materials:** Do you wish to include the location of the underlying research materials (e.g. data, samples or models) for your article? If so, please insert this sentence before the reference section: ‘The underlying research materials for this article can be accessed at <full link>/ description of location [author to complete]’. If your article includes supplemental data, the link will also be provided in this paragraph. See <<http://journalauthors.tandf.co.uk/preparation/multimedia.asp>> for further explanation of supplemental data and underlying research materials.
6. The **CrossRef database** (www.crossref.org/) has been used to validate the references. Changes resulting from mismatches are tracked in **red** font.

AQ1	Please suggest whether the term “Dicarbxylic” could be changed to “Dicarboxylic” in the article title. And also suggest whether serial comma can be given in the article title.
AQ2	Please note that the references have been renumbered in the text as well as in the list so as to maintain sequential order. Kindly check and confirm the change
AQ3	Please check whether the equations have been correctly formatted throughout the article
AQ4	Please provide the Notes on contributors (regarding current academic position, research interests, books/articles published).
AQ5	The disclosure statement has been inserted. Please correct if this is inaccurate.
AQ6	Please provide an English translation of the title in Ref. [6].
AQ7	Please provide an English translation of the title in Ref. [8].
AQ8	Please provide missing publisher for Ref. [16].

How to make corrections to your proofs using Adobe Acrobat/Reader

Taylor & Francis offers you a choice of options to help you make corrections to your proofs. Your PDF proof file has been enabled so that you can mark up the proof directly using Adobe Acrobat/Reader. This is the simplest and best way for you to ensure that your corrections will be incorporated. If you wish to do this, please follow these instructions:

1. Save the file to your hard disk.
2. Check which version of Adobe Acrobat/Reader you have on your computer. You can do this by clicking on the “Help” tab, and then “About”.

If Adobe Reader is not installed, you can get the latest version free from <http://get.adobe.com/reader/>.

3. If you have Adobe Acrobat/Reader 10 or a later version, click on the “Comment” link at the right-hand side to view the Comments pane.
4. You can then select any text and mark it up for deletion or replacement, or insert new text as needed. Please note that these will clearly be displayed in the Comments pane and secondary annotation is not needed to draw attention to your corrections. If you need to include new sections of text, it is also possible to add a comment to the proofs. To do this, use the Sticky Note tool in the task bar. Please also see our FAQs here: <http://journalauthors.tandf.co.uk/production/index.asp>.
5. Make sure that you save the file when you close the document before uploading it to CATS using the “Upload File” button on the online correction form. If you have more than one file, please zip them together and then upload the zip file.

If you prefer, you can make your corrections using the CATS online correction form.

Troubleshooting

Acrobat help: <http://helpx.adobe.com/acrobat.html>

Reader help: <http://helpx.adobe.com/reader.html>

Please note that full user guides for earlier versions of these programs are available from the Adobe Help pages by clicking on the link “Previous versions” under the “Help and tutorials” heading from the relevant link above. Commenting functionality is available from Adobe Reader 8.0 onwards and from Adobe Acrobat 7.0 onwards.

Firefox users: Firefox’s inbuilt PDF Viewer is set to the default; please see the following for instructions on how to use this and download the PDF to your hard drive: http://support.mozilla.org/en-US/kb/view-pdf-files-firefox-without-downloading-them#w_using-a-pdf-reader-plugin

Mixed-ligand complex formation equilibria of vanadium(III) with picolinic and dipicolinic acids with some dicarboxylic acids (oxalic, malonic and phthalic acids) studied in 3.0 M KCl at 25 °C

5 Ana Esteves^a, Jessaid Escobar^a, José Daniel Martínez^a, Mary Lorena Araujo^a, Felipe Brito^a, Lino Hernández^b,
Edgar Del Carpio^b and Vito Lubes^{b*}

^aFacultad de Ciencias, Escuela de Química, Centro de Equilibrios en Solución, Universidad Central de Venezuela (UCV), Caracas, Venezuela; ^bDepartamento de Química, Universidad Simón Bolívar (USB), Caracas, Venezuela

10 The ternary complexes formed between vanadium(III) and picolinic acid (Hpic) and dipicolinic acid (H₂dipic) with the
dicarboxylic oxalic, malonic, and phthalic acids (H₂ox), (H₂mal), and (H₂phtha) briefly (H₂L) were studied in aqueous
solution by means of electromotive forces' measurements emf(H) at 25 °C and 3.0 mol dm⁻³ KCl as ionic medium. The
potentiometric data were analyzed using the least-squares computational program LETAGROP, considering the hydrolysis
15 products of the V(III), the acidity constants of the ligands employed, and the formation constants of the binary complexes,
obtaining the complexes V(pic)(ox), log β_{1,1,1,-3} = 5.39(7) and [V(pic)₂(ox)]⁻, log β_{1,2,1,-4} = 4.3(1); V(pic)(ma),
log β_{1,1,1,-3} = 5.58(5) and [V(pic)₂(ox)]⁻, log β_{1,2,1,-4} = 5.00(8); V(pic)(phtha), log β_{1,1,1,-3} = 5.66(6) and [V(pic)₂(phtha)]⁻
log β_{1,2,1,-4} = 4.5(2) in the vanadium(III)-Hpic-H₂L system. In the case of the vanadium(III)-H₂dipic-H₂L system,
the complexes [V(dipic)(ox)]⁻, log β_{1,1,1,-4} = 9.3(2), [V(dipic)(Hox)(ox)]²⁻, log β_{1,1,2,-5} = 11.2(2) and [V(dipic)(ox)₂]³⁻,
log β_{1,1,2,-6} = 5.7 max 7.0; [V(dipic)(ma)]⁻, log β_{1,1,1,-4} = 8.7(2), [V(dipic)(Hma)(ma)]²⁻, log β_{1,1,2,-5} = 10.3(1) and [V
20 (dipic)(ma)₂]³⁻, log β_{1,1,2,-6} = 6.9(1); [V(dipic)(phtha)]⁻, log β_{1,1,1,-4} = 10.9(1), [V(dipic)(Hphtha)(phtha)]²⁻, log
β_{1,1,2,-5} = 9.2 max 10.4 and [V(dipic)(phtha)₂]³⁻, log β_{1,1,2,-6} = 6.9(3) were observed.

Keywords: vanadium(III) complexes; picolinic acid; dipicolinic acid; dicarboxylic acids; potentiometric studies; speciation; ternary complexes

1. Introduction

25 Recent studies on the coordination chemistry of vanadium have shown that V complexes have great potential
as pharmaceuticals. Studies have been conducted *in vivo* and *in vitro* of these compounds in different oxidation
states, and the complete structure of some has been determined. However, there are many vanadium complexes
30 that have not been isolated and studied structurally.[1]

Among the most common biochemical properties that support the study of the coordination chemistry of vanadium
are the metabolic enzyme inhibition of ATPase: stimulation of phosphomutases; a required cofactor for a
35 number of haloperoxidases; regulation of anomalies of thyroid; insulinomimetic and anticarcinogenic activity;
and in some cases a preventive and therapeutic capacity of the malignancy of certain tumor cell lines.[1,2]

40 The interest in the study of the aqueous chemistry of vanadium began with the discovery of vanadium in some
species of tunicates. Even when high concentrations of vanadium have been found frequently in various tissues
of living organisms, few animal organisms are known to accumulate vanadium. However, high levels of vanadium
45 have been reported in ascidians (*Pseudopotamilla acce-lata*), discovered by German M. Henze in 1911 and is so
far the second tunicate with high accumulation in the

animal kingdom.[3] At the bottom of the Black Sea bay located in Sevastopol, about 400 kg of vanadium is
50 found per square mile, located in the blood and tissues of these sedentary tunicates.[4] The species *Ascidia gem-
mata* contains values as high as 0.35 M of vanadium in the blood cells corresponding to 107 times the concentra-
tion of the metal in sea water.[5]

55 In this work, we report the results on mixed-ligand complex formation in the V(III)-picolinic acid-H₂L and
V(III)-dipicolinic acid-H₂L systems, where the H₂L were the dicarboxylic oxalic, malonic, and phthalic acids
(H₂ox), (H₂mal), and (H₂phtha). The pH-potentiometry technique was used to determine the stoichiometries and
60 stability constants of the complexes formed in aqueous solution, because the study of vanadium(III) complexes
gives you an idea of the behavior of these complexes in natural systems

2. Experimental

2.1. Reagents


70 The VCl₃ (Merck p.a); and the pyridinecarboxylic acids, picolinic acid (Hpic), and dipicolinic acid (H₂dipic)
(Merck p.a.); and the dicarboxylic oxalic, malonic and phthalic acids (H₂ox), (H₂mal), and (H₂phtha) briefly
(H₂L) also (Merck p.a) were used without purification.

*Corresponding author. Email: lubesv@usb.ve

2

A. Esteves et al.

The HCl and KOH solutions were prepared using 100.0 mmol dm⁻³ Titrisol Merck ampoules. The KOH solution was standardized against potassium hydrogen phthalate. The solutions were prepared using triple glass-distilled water and boiled before the preparation of the solutions in order to remove dissolved CO₂. To prevent the hydrolysis of the VCl₃ stock solution, it contained 200 mmol dm⁻³ HCl and was maintained under a H₂ atmosphere in the presence of a Pt platinized net in order to avoid oxidation of the V(III) solution to V(IV).[6] In this case, the H₂ cannot induce reduction to V(II), because the V(III)/V(II) standard potential is negative ($E^0 = -0.26$ V).[7] Moreover, if there is any oxidation to V(IV), the solution is immediately reduced to V(III) because of the standard potential of the reaction,





which is 366.3(3) mV.[8] Under these conditions, the V(III) solution can be maintained. The stability of the V(III) solution was checked periodically by spectrophotometric measurements, and it was shown to be stable for several weeks. The emf (H) measurements were carried out in aqueous solution at an ionic strength of 3.0 mol dm⁻³ in KCl. Nitrogen-free O₂ and CO₂ was used to maintain an inert atmosphere.

2.2. Methods

The emf (H) measurements were done using the following instruments: Thermo Orion model 520A pH meter, Metrohm EA 876-20 titration vessel, Lauda Brikmann RM6 thermostat bath, Shimadzu UV-1601 PC spectrophotometer, and a quartz cell with a 10.0-mm path length. The sealed 100-mL thermostatted double-walled glass titration vessel was fitted with a combined Orion Ross 8102BN pH electrode with a titrant inlet, magnetic stirrer, and an inert nitrogen atmosphere inlet with outlet tubes. The temperature was maintained at 25.0(1) °C by constant circulation of water from the thermostat bath.

The emf (H) measurements were carried out by means of the REF//S/GE cell, where REF = Ag, AgCl/3.0 mol dm⁻³ KCl; S = equilibrium solution; and GE = glass electrode. At 25 °C, the emf (mV) of this cell follows the Nernst equation, $E = E^0 + jh + 59.16 \log h$, where h represents the free hydrogen ion concentration, E^0 is the standard potential, and j is a constant which takes into account the liquid junction potential.[9] The experiments were carried out as follows: a fixed volume of 0.100 mol dm⁻³ HCl was titrated with successive additions of 0.100 mol dm⁻³ KOH until near neutrality in order to get the parameters E^0 and j . Then, aliquots of 6Mepic, and the ligands B and an aliquot of the Vanadium(III) stock solution were added sequentially. Finally, the titration was continued with 0.100 mol dm⁻³ KOH. The measurements were done using a total metal concentration, $M_T = 2-3$ mmol dm⁻³ and Vanadium(III): Hpic: H₂L and Vanadium(III): H₂dipic: H₂L molar ratios $R = 1:1:1, 1:2:1, \text{ and } 1:1:2$.

The systems Vanadium(III): Hpic: H₂L and Vanadium(III): H₂dipic: H₂L were studied according to the reactions scheme:



where H₂L represents the ligands: H₂ox, H₂mal, and H₂phtha, $[\text{V}q(\text{OH})p(\text{Hpic})r(\text{H}_2\text{L})s]$ and $[\text{V}q(\text{OH})p(\text{H}_2\text{dipic})r(\text{H}_2\text{L})s]$ are the ternary (p, q, r, s) complexes, and $\beta_{p,q,r,s}$ are the respective stability constant.

The potentiometric data were analyzed using the program LETAGROP,[10,11] in order to minimize the function $Z_B = (h - H)/M_T$, where H is the total (analytical) concentration of H⁺, h represents the equilibrium concentration of H⁺, and M_T represents the total (analytical) concentration of V³⁺. Equilibria corresponding to the formation of the hydroxo complexes of V(III) were considered in the calculation of the stability constants of the ternary complexes. The following species were assumed: $[\text{V}(\text{OH})]^{2+}$, $\log \beta_{1,-1} = -3.07(3)$; $[\text{V}_2\text{O}]^{4+}$, $\log \beta_{2,-2} = -3.94(2)$; and $[\text{V}_2\text{O}(\text{OH})]^{3+}$, $\log \beta_{2,-3} = -7.87(9)$. [8] In the Binary V(III)-Hpic [12] system were considered the complexes: $[\text{V}(\text{pic})]^{2+}$, $\log \beta_{1,1,-1} = 0.39(9)$; $[\text{V}(\text{pic})(\text{OH})]^{+}$, $\log \beta_{1,1,-2} = -2.12(7)$; $[\text{V}(\text{pic})_2]$, $\log \beta_{1,1,-2} = 1.68(2)$; $\text{V}(\text{pic})_3$, $\log \beta_{1,1,-3} = 0.68(3)$; $\text{V}_2\text{O}(\text{pic})_4$, $\log \beta_{2,4,-6} = -0.55(4)$; In the V(III)-H₂dipic [12] system were considered the complexes: $[\text{V}(\text{dipic})]^{+}$, $\log \beta_{1,1,-2} = 1.16(2)$; $\text{V}(\text{dipic})(\text{OH})$, $\log \beta_{1,1,-3} = -3.38(3)$; $[\text{V}(\text{dipic})(\text{OH})_2]^{-}$, $\log \beta_{1,1,-4} = -10.36(2)$; $\text{V}(\text{Hdipic})(\text{dipic})$, $\log \beta_{1,2,-3} = 1.85(3)$; $[\text{V}(\text{dipic})_2]^{-}$, $\log \beta_{1,2,-4} = -0.96(2)$. In the V(III)-H₂ox [13] system were the complexes: $[\text{V}(\text{Hox})]^{2+}$, $\log \beta_{1,1,-1} = 6.83(3)$; $[\text{V}(\text{ox})]^{2+}$, $\log \beta_{1,1,-2} = 5.16(3)$; $[\text{V}(\text{ox})(\text{OH})]$, $\log \beta_{1,1,-3} = 0.97(5)$; $[\text{V}(\text{ox})(\text{OH})_2]^{-}$, $\log \beta_{1,1,-4} = -4.76(7)$; $[\text{V}(\text{ox})_2]^{-}$, $\log \beta_{1,2,-4} = 6.37(8)$; $[\text{V}(\text{ox})_2(\text{OH})]^{2-}$, $\log \beta_{1,2,-5} = 1.6(1)$; $[\text{V}(\text{ox})_3]^{3-}$, $\log \beta_{1,3,-6} = 7.23(6)$; $[\text{V}(\text{ox})_3(\text{OH})]^{4-}$, $\log \beta_{1,3,-7} = 1.15(7)$. In the V(III)-H₂mal [13] system were the complexes: $[\text{V}(\text{Hmal})]^{2+}$, $\log \beta_{1,1,-1} = 6.74(3)$; $[\text{V}(\text{mal})]^{2+}$, $\log \beta_{1,1,-2} = 5.04(3)$; $[\text{V}(\text{mal})_2]^{-}$, $\log \beta_{1,2,-4} = 7.11(7)$; $[\text{V}(\text{mal})_2(\text{OH})]^{2-}$, $\log \beta_{1,2,-5} = 4.07(5)$; $[\text{V}(\text{mal})_2(\text{OH})_2]^{-}$, $\log \beta_{1,2,-6} = -1.05(8)$; $[\text{V}(\text{mal})_3]^{3-}$, $\log \beta_{1,3,-6} = 4.09(6)$; $[\text{V}(\text{ox})_3(\text{OH})]^{4-}$, $\log \beta_{1,3,-7} = 0.34(7)$. In the V(III)-H₂phtha [14] system were considered the complexes: $[\text{V}(\text{Hphtha})]^{2+}$, $\log \beta_{1,1,-1} = 7.84(5)$; $[\text{V}(\text{phtha})]^{+}$, $\log \beta_{1,1,-2} = 4.9(1)$; $[\text{V}(\text{phtha})_2]^{-}$, $\log \beta_{1,2,-4} = 8.04(4)$; $\text{V}(\text{OH})(\text{phtha})$, $\log \beta_{1,1,-3} = 1.55(8)$; $\text{V}(\text{Hphtha})(\text{phtha})_2$, $\log \beta_{1,2,-3} = 8.5(1)$; $[\text{V}(\text{OH})(\text{phtha})_2]^{2-}$, $\log \beta_{1,2,-5} = 4.2(2)$; $[\text{V}(\text{OH})_2(\text{phtha})_2]^{2-}$, $\log \beta_{1,2,-6} = 0.8(1)$; $[\text{V}(\text{phtha})_3]^{3-}$, $\log \beta_{1,3,-6} = 4.88(5)$.

The stability constant of the V(III) hydroxo complexes, the stability constants of the ligands, and the stability constants of binary complexes were kept fixed

60

65

70

75

80

85

90

95

100

105

110

during the analysis. The aim of the analysis was to find a complex or complexes giving the lowest sum of the errors squared, $U = \sum (Z_B^{\text{exp}} - Z_B^{\text{calc}})^2$, the fittings were done by testing different (p, q, r, s) combinations.

The species distribution diagrams were carried out using the computer program HYSS,[15] yielding the $\beta_{p,q,r,s}$ values, which are summarized in Tables 2 and 3.

3. Results and discussion

3.1. Ionization constants of ligands studied

The analysis of the picolinic acid (Hpic) system was obtained two pK_a values, being the first pK_a value that corresponds to the deprotonation of carboxylic group, and the second pK_a corresponds to the deprotonation of the pyridinic group. In the case of the dipicolinic acid ($H_2\text{dipic}$) system, two pK_a values were also obtained, the first pK_a corresponds to the deprotonation of the first carboxylic group, which has a value of 0.5 and it is not possible to measure it due to the high error made with the glass electrode. The second pK_a value corresponds to the deprotonation of the second carboxylic group, and the last pK_a corresponds to the deprotonation of the pyridinic group.[12] In the case of the dicarboxylic oxalic, malonic, and phthalic acids ($H_2\text{ox}$), ($H_2\text{mal}$), and ($H_2\text{phtha}$) briefly (H_2L), they have two pK_a that correspond to the deprotonation of the carboxylic groups. The pK_a values obtained are in good agreement with the values reported in the literature [16,17] (Table 1).

3.2. Ternary vanadium(III) complexes

3.2.1. V(III)-Hpic- H_2L system

Table 2 presents the results of the analysis performed with the LETAGROP program in which it were observed the formation of the complexes $V(\text{pic})(L)$ and $[V(\text{pic})_2(L)]^-$. The respective species distribution diagrams are given in Figures 13.

3.2.2. V(III)- $H_2\text{dipic}$ - H_2L system

The results of the analysis are shown in Table 3 and were observed the complexes $[V(\text{dipic})(L)]^-$, $[V(\text{dipic})(HL)(L)]^{2-}$, and $[V(\text{dipic})(L)_2]^{3-}$ in all the systems. The stability constants given and the respective species distribution diagrams are presented in Figures 4–6.

In the V(III)-pic- H_2L ternary system, two complexes were detected: $V(\text{pic})(L)$ and $[V(\text{pic})_2(L)]^-$. In Figure 1 shows the species distribution diagram for the V(III)-pic- $H_2\text{ox}$ system, taking into account the molar ratio $V^{3+}:\text{Hpic}:\text{H}_2\text{ox}$ $R = 1:2:1$ and $M_T = 3 \text{ mmol dm}^{-3}$. The results show that the ternary complex is more abundant, $V(\text{pic})(L)$ predominates over a wide pH range ($\text{pH} \leq 1 \leq 7$), while the complex $[V(\text{pic})_2(L)]^-$ is in low proportion. In the acid region, the binary complex $[VL]^{2-}$ dominates in the range $1 \leq \text{pH} \leq 4$. At $\text{pH} > 6.5$, the dimeric complex V_2OL_4 predominates. Figure 2 shows the species distribution diagrams for the V(III)-pic- $H_2\text{mal}$ system taking into account the molar ratio $V^{3+}:\text{Hpic}:\text{H}_2\text{mal}$ $R = 1:2:1$ and $M_T = 3 \text{ mmol dm}^{-3}$. This system has the same

Table 1. Ligands acidity constants ($\log \beta_{p,r}$) in 3.0 M KCl at 25 °C.

Equilibrium	Hpic	$\log \beta_{p,r}$ $H_2\text{dipic}$	$H_2\text{ox}$	$H_2\text{mal}$	$H_2\text{phtha}$
$HL + H^+ \rightleftharpoons H_2L^+$	1.51(2)				
$H_nL \rightleftharpoons H_{n-1}L^- + H^+$	-5.57(2)	2.45(2)	-0.68(3)	-2.87(3)	-2.95(3)
$H_nL \rightleftharpoons H_{n-2}L^{2-} + 2H^+$		-7.11(3)	-4.47(3)	-8.15(5)	-7.71(4)
Dispersion σ (Z)	0.020	0.026	0.022	0.026	0.021
Ligand concentration (mmol dm^{-3})	3.0	3.0	3.0	3.0	3.0
pK_i					
$H_2L^+ - HL$	1.51	2.45			
$HL - L^-$	5.57	4.66			
$H_2L - HL$			0.68	2.87	2.95
$HL - L^-$			3.79	5.28	4.76

Note: Values in parentheses are standard deviations [$3\sigma(\log \beta)$] on the last significant figure.

Table 2. Stability constants ($\log \beta_{p,q,r,s}$) of the V(III)-Hpic- H_2L systems (KCl 3.0 mol dm^{-3} , 25 °C). Considering the following reaction scheme: $pH_2O + qV^{3+} + rHpic + sH_2L \rightleftharpoons [Vq(OH)p(Hpic)r(H_2L)s] + pH^+$.

Complex	Model (p, q, r, s)	Oxalic acid	Malonic acid	Phthalic acid
$V(\text{pic})L$	(-3, 1, 1, 1)	5.39 (7)	5.58 (5)	5.66 (6)
$[V(\text{pic})_2L]^-$	(-4, 1, 2, 1)	4.3 (1)	5.00 (8)	4.5 (2)
Dispersion (σ)		0.089	0.063	0.088

Note: Values in parentheses are standard deviations [$3\sigma(\log \beta_{p,q,r,s})$] on the last significant figure.

4

A. Esteves et al.




Figure 1. Species distribution diagram of the V(III)-Hpic-H₂Ox system. Considering $M_T = 3$ mM and V^{3+} -Hpic-H₂Ox ratio $R = 1:2:1$.

Table 3. Stability constants ($\log \beta_{p,q,r,s}$) of the V(III)-H₂dipic-H₂L systems (KCl 3.0 mol dm⁻³, 25 °C). Considering the following reaction scheme: $pH_2O + qV^{3+} + rH_2dipic + sH_2L \rightleftharpoons [Vq(OH)p(H_2dipic)r(H_2L)s] + pH^+$.

Complex	(<i>p, q, r, s</i>)	Oxalic acid	$\log \beta_{p,q,r,s}$ Malonic acid	Phthalic acid
[V(dipic)L] ⁻	(-4, 1, 1, 1)	9.3 (2)	8.7 (2)	10.9 (1)
[V(dipic)(HL)L] ²⁻	(-5, 1, 1, 2)	11.2 (2)	10.3 (1)	9.2 max 10.4
[V(dipic)L ₂] ³⁻	(-6, 1, 1, 2)	5.7 max 7.0	6.9 (1)	6.9 (3)
Dispersion (σ)		0.073	0.132	0.096

Note: Values in parentheses are standard deviations [$3\sigma(\log \beta_{p,q,r,s})$] on the last significant figure.




Figure 2. Species distribution diagram of the V(III)-Hpic-H₂mal system. Considering $M_T = 3$ mM and V^{3+} -Hpic-H₂mal ratio $R = 1:2:1$.

5 behavior as the V(III)-pic-H₂Ox, but in this case, the
 complexes V(pic)(L) and [V(pic)₂(L)]⁻ are formed in a
 higher proportion and the dimeric complex V₂OL₄ is
 formed in less proportion. In Figure 3 is given the spe-
 10 cies distribution diagram for the V(III)-pic-H₂phtha
 system taking into account the molar ratio V^{3+} :Hpic:H₂mal
 $R = 1:2:1$ and $M_T = 3$ mmol dm⁻³, but in this case, the
 complex V(pic)(L) is the predominant species in the pH
 range 2–3.8, the [V(pic)₂(L)]⁻ in formed in low

quantities and the species V₂OL₄, [V(OH)₂L₂]³⁻, and
 [VL₃]³⁻ predominates at pH > 4.4.

In the V(III)-dipic-H₂L ternary system, three com-
 15 plexes were detected: [V(dipic)(L)]⁻, [V(dipic)(HL)
 (L)]²⁻, and [V(dipic)(L)₂]³⁻. Figure 4 shows the species
 distribution diagrams for the V(III)-H₂dipic-H₂Ox system
 20 under the experimental conditions: molar ratio V^{3+} :
 H₂dipic:H₂Ox $R = 1:1:2$ and $M_T = 3$ mmol dm⁻³ and
 shows that the complex [V(dipic)L]⁻ is present in the




Figure 3. Species distribution diagram of the V(III)-Hpic-H₂phtha system. Considering $M_T = 3$ mM and V^{3+} -Hpic-H₂phtha ratio $R = 1:2:1$.




Figure 4. Species distribution diagram of the V(III)-H₂dipic-H₂ox system. Considering $M_T = 3$ mM and V^{3+} -H₂dipic-H₂ox ratio $R = 1:1:2$.




Figure 5. Species distribution diagram of the V(III)-H₂dipic-H₂mal system. Considering $M_T = 3$ mM and V^{3+} -H₂dipic-H₂mal ratio $R = 1:1:2$.

pH range 1–7, and in less proportion, it is formed the acidic complex $[V(dipic)(HL)L]^{2-}$ in the range $1 < \text{pH} < 4$, and the complexes $[VL_3]^{3-}$ and $[V(OH)L_3]^{4-}$ are formed in low proportions. The species distribution diagrams for the system V(III)-H₂dipic-H₂mal are given in Figure 5 for the conditions molar ratio V^{3+} :

H₂dipic:H₂mal $R = 1:1:2$ and $M_T = 3$ mmol dm⁻³, show the formation in less than 50% of the $[V(dipic)(L)]^-$ complex in the pH range 1–4, the complex $[V(dipic)(HL)(L)]^{2-}$ predominates between 1.2 and 3.3 pH range,




Figure 6. Species distribution diagram of the V(III)-H₂dipic-H₂phtha system. Considering $M_T = 3$ mM and V^{3+} -H₂dipic-H₂phtha ratio $R = 1:1:2$.

6

A. Esteves et al.

and the $[V(\text{dipic})(L)_2]^{3-}$ complex is the most important species at $\text{pH} > 3.2$. Finally, Figure 6 shows the species distribution diagrams for the $V(\text{III})\text{-H}_2\text{dipic-H}_2\text{phtha}$ system taking into account the conditions molar ratio $V^{3+}:\text{H}_2\text{dipic}:\text{H}_2\text{mal}$ $R = 1:1:2$ and $M_T = 3 \text{ mmol dm}^{-3}$, the $[V(\text{dipic})(L)]^-$ complex is very important in the pH range 1–4 and is formed in less extension the complex the $[V(\text{dipic})(L)_2]^{3-}$ and the most important species at $\text{pH} > 5$ is the complex $[V(\text{OH})_2(L)_2]^{3-}$.

4. Conclusions

In this work, we have studied the speciation of the ternary complexes of Vanadium(III)-Hpic and Vanadium(III)- H_2dipic with the dicarboxylic oxalic, malonic, and phthalic acids (H_2ox), (H_2mal) and (H_2phtha). The data analyzed using the least-squares computational program LETAGROP indicate the formation of the complexes $V(\text{pic})(L)$ and $[V(\text{pic})_2(L)]^-$ in the vanadium(III)-Hpic- H_2L system. In the case of the vanadium(III)- $\text{H}_2\text{dipic-H}_2L$ system were observed the complexes $[V(\text{dipic})(L)]^-$, $[V(\text{dipic})(\text{HL})(L)]^{2-}$ and $[V(\text{dipic})(L)_2]^{3-}$.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The authors are grateful for financing provided by the Decanato de Investigación y Desarrollo, from the Simón Bolívar University. Lino Hernández and Edgar del Carpio thanks scholarship from Simón Bolívar University.

References

- [1] Da Costa F, Lubes G, Rodríguez M, Lubes V. Study of the ternary complex formation between vanadium(III), dipicolinic acid and small blood serum bioligands. *J Solution Chem.* 2011;40:106–117.
- [2] Osińska-Królicka I, Podsiadły H, Bukietyńska K, Zemanek-Zboch M, Nowak D, Suchoszek-Lukaniuk K, Malicka-Błaszkiwicz M. Vanadium(III) complexes with l-cysteine – stability, speciation and the effect on actin in hepatoma Morris 5123 cells. *J. Inorg. Biochem.* 2004;98:2087–2098.

- [3] Crans DC, Smeey JJ, Gaidamauskas E, Yang L. The chemistry and biochemistry of vanadium and the biological activities exerted by vanadium compounds. *Chem. Rev.* 2004;104:849–902.
- [4] Senozan NM. Vanadium in the living world. *J. Chem. Edu.* 1974;51:503–505.
- [5] Kanamori K. Structures and properties of multinuclear vanadium(III) complexes: seeking a clue to understand the role of vanadium(III) in ascidians. *Coord. Chem. Rev.* 2003;237:147–161.
- [6] Mateo S, Brito F. Complejos de vanadio(III). 1. Hidrólisis de V^{3+} en $(K)Cl$ 3.0 M a 25 °C. *An. Fis. Quím.* 1968;64B:115–138. [AQ6](#)
- [7] Lannon AM, Lappin AG, Segal MG. Electron-transfer reactions of tris(picolinato)vanadate(II), a LOMI reagent. *Inorg. Chem.* 1984;23:4167–4170. 55
- [8] Brito F, Gonçalves JM. Complejos de Vanadio(III) 2. Potencial Normal del Sistema VO^{2+}/V^{3+} [25 °C $(K)Cl$ 3 M]. *An. Fis. Quím.* 1982;78:104–105. [AQ7](#)
- [9] Biedermann G, Sillén LS. The hydrolysis of metal ions. IV. Liquid-junction potentials and constancy of activity factors in $NaClO_4\text{-}HClO_4$ ionic medium. *Ark. Kemi.* 1953;5:425–540. 60
- [10] Brito F, Araujo ML, Lubes V, D’Ascoli A, Mederos A, Gili P, Domínguez S, China E, Hernández-Molina R, Armas MT, Baran EJ. Emf(H) data analysis of weak metallic complexes using reduced formation functions. *J. Coord. Chem.* 2005;58:501–512. 65
- [11] Sillén LG, Wamqvist B. High-speed computers as a supplement to graphical methods. VI. A strategy for two-level Letagrop adjustment of common and “group” parameters. Features that avoid divergence. *Arkiv Kemi.* 1969;31:315–339. 70
- [12] Lubes V. Vanadium(III) complexes with picolinic acid and dipicolinic acid in aqueous solution. *J. Solution Chem.* 2005;34:899–915. 75
- [13] Bricual J, Lubes V, Araujo ML, Brito F. Vanadium(III) complexes in aqueous solution with the dicarboxylic oxalic, malonic and succinic acids. *J. Chill. Chem. Soc.* 2004;49:285–288. 80
- [14] Gonçalves M, Martínez JD, Araujo ML, Brito F, Lubes G, Rodríguez M, Lubes V. Speciation of the vanadium (III) complexes with 6-methylpicolinic acid, salicylic acid and phthalic acid. *J Solution Chem.* 2011;40:1517–1527.
- [15] Alderighi L, Gans P, Ienco A, Peters D, Sabatini A, Vacca A. Hyperquad simulation and speciation (HySS): a utility program for the investigation of equilibria involving soluble and partially soluble species. *Coord. Chem. Rev.* 1999;184:311–318. 85
- [16] Martell AE, Smith RM, Motekaitis RJ. NIST standard reference database. Version 8. Gaithersburg, MD; 2004. [AQ80](#)
- [17] Pettit D, Powell K. IUPAC stability constants database. Otley (UK): Academic Software; 1997.